

16-MAR-2007

26-AUG-2008 Notification - Telephone Number

Safer's

BTK™

Biological Insecticide

Controls caterpillars, including cabbage worm, tomato hornworm, tent caterpillars, gypsy moth, leafrollers and other listed insects on vegetables, fruits, shade trees, ornamentals and evergreens.

Gypsy moth

Tent Caterpillar

Cabbage Looper

GUARANTEE: Bacillus thuringiensis sub-species kurstaki.

Potency: 10,600 International Units per mg,

Equivalent to 12.7 Billion International Units (BIU) per litre.

Warning, contains the allergen sulfites

REG.NO. 24536 P.C.P. ACT

READ THE LABEL BEFORE USING

DOMESTIC

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS 100 mL

Concentrate

Notification Change

Woodstream Canada Corporation, 4 Lowry Drive, Brampton, ON L7A 1C4 1-800-800-1819

Safer's BTK™ Biological Insecticide
SHAKE WELL BEFORE USING.

APPLICATION RATE: Add 30 mL of concentrate to 10 litres of water. 10 L of spray solution will cover approximately 100 square metres. Spray all leaf surfaces thoroughly to provide good coverage without run-off. Repeat, if required, every 7-10 days.

DIRECTIONS FOR USE: Safer's BTK™ Biological Insecticide is highly specific to some kinds of caterpillars. Treat caterpillars when they are young to obtain best results. Safer's BTK™ Biological Insecticide is a stomach poison and therefore must be ingested by the caterpillar to be effective. After ingestion of a sufficient product dose, feeding stops immediately and death will follow in 2-5 days.

FOR EXPANDED DIRECTIONS FOR USE, SEE INSIDE LABEL.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. STORE IN COOL PLACE AT TEMPERATURES BETWEEN 0° C AND 20° C.

When stored under optimum conditions, (15° C or less), this product should be used within 2 years of the date of manufacture. Avoid contact with skin, eyes and clothing. Avoid direct application to ponds, lakes and streams. Do not contaminate water bodies by cleaning equipment or disposal of waste. Avoid application if rainfall is imminent.

DISPOSAL: Do not reuse empty container. Wrap empty container and dispose in household garbage.

FIRST AID: If swallowed, drink 1-2 glasses of water. If in eyes, flush with water. If on skin, wash with soapy water. If irritation persists, consult a physician. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

(Inside Label)

SHAKE WELL BEFORE USING.

DIRECTIONS FOR USE: Safer's BTK™ Biological Insecticide is highly specific to some kinds of caterpillars. Treat caterpillars when they first appear and are actively feeding for best results. Safer's BTK™ Biological Insecticide is most effective when caterpillars are small and young. This product is not effective on adult moths. Safer's BTK™ Biological Insecticide is a stomach poison and therefore must be ingested by the caterpillar to be effective. After ingestion of a sufficient product dose, feeding stops immediately and death will follow in 2-5 days. Add some water to the sprayer and then add the correct amount of Safer's BTK™ Biological Insecticide. Continue filling the sprayer with water until the correct amount of spray solution is produced. Shake sprayer periodically during use to ensure product is properly mixed. Use within 12 hours of mixing. May be applied through any standard back-pack or hand-held sprayer.

APPLICATION RATE: Add 30 mL of concentrate to 10 litres of water. 10 L of spray solution will cover approximately 100 square metres. Spray all leaf surfaces thoroughly to provide good coverage without run-off. Repeat, if required, 7-10 days later.

VEGETABLES AND FRUITS: On Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kale, Lettuce, Mustard Greens, Spinach and Tomatoes for control of Cabbage Looper, Imported Cabbageworm and Tomato Hornworm. In greenhouse plantings of Cucumbers, Peppers and Tomatoes for control of Cabbage Looper. On Raspberries, and on Pears and Apples for control of Leafrollers. Mix 30 mL of Safer's BTK™ Biological Insecticide concentrate in 10 litres of water

and spray all leaf surfaces thoroughly to provide good coverage without run-off at first sign of infestation. Repeat at 7-10 day intervals, if required.

SHADE TREES AND ORNAMENTALS: On Shade trees and Ornamentals for control of Spring and Fall Cankerworm, Tent Caterpillar, Satin Moth, and Gypsy Moth. Mix 30 mL of Safer's BTK™ Biological Insecticide concentrate in 10 litres of water and spray all leaf surfaces thoroughly to provide good coverage without run-off when leaf expansion reaches 40-50% and caterpillars first appear. Repeat in 7-10 days, if required.

EVERGREENS: On Evergreens for control of Spruce budworm, Jack Pine budworm and Eastern Hemlock Looper. Mix 30 mL of Safer's BTK™ Biological Insecticide concentrate in 10 litres of water and spray all leaf surfaces thoroughly to provide good coverage without run-off when caterpillars first appear. Repeat in 7-10 days, if required.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. STORE IN A COOL PLACE AT TEMPERATURES BETWEEN 0° C and 20° C. When stored under optimum conditions, (15° C or less), this product should be used within 2 years of the date of manufacture. Avoid contact with skin, eyes, and clothing. Avoid direct application to ponds, lakes or streams. Do not contaminate water bodies by cleaning equipment or disposal of waste. Avoid application if rainfall is imminent.

DISPOSAL: Do not reuse empty container. Wrap empty container and dispose of in household garbage.

FIRST AID: If swallowed, drink 1-2 glasses of water. If in eyes, flush with water. If on skin, wash with soapy water. If irritation persists, consult of physician. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

QUESTIONS? Call Woodstream Canada Corporation at 1-800-800-1819.

Notification Change

Safer's BTK™ Biological Insecticide

SHAKE WELL BEFORE USING.

APPLICATION RATE: Add 30 mL of concentrate to 10 litres of water. 10 L of spray solution will cover approximately 100 square metres. Spray all leaf surfaces thoroughly to provide good coverage without run-off. Repeat, if required, every 7-10 days.

DIRECTIONS FOR USE: Safer's BTK™ Biological Insecticide is highly specific to some kinds of caterpillars. Treat caterpillars when they are young to obtain best results. Safer's BTK™ Biological Insecticide is a stomach poison and therefore must be ingested by the caterpillar to be effective. After ingestion of a sufficient product dose, feeding stops immediately and death will follow in 2-5 days.

FOR EXPANDED DIRECTIONS FOR USE, SEE INSIDE LABEL.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. STORE IN A COOL PLACE AT TEMPERATURES BETWEEN 0° C and 20° C. When stored under optimum conditions, (15° C or less), this product should be used within 2 years of the date of manufacture. Avoid contact with skin, eyes, and clothing. Avoid direct application to ponds, lakes or streams. Do not contaminate water bodies by cleaning equipment or disposal of waste. Avoid application if rainfall is imminent.

DISPOSAL: Do not reuse empty container. Wrap empty container and dispose of in household garbage.

FIRST AID: If swallowed, drink 1-2 glasses of water. If in eyes, flush with water. If on skin, wash with soapy water. If irritation persists, consult of physician. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.